

63-3-4

AFCRC TN AFCRL - 63 - 288

ASTIA Document No.

THE STRENGTHS OF ABSORPTION LINES IN THE 2-0 BAND
OF CARBON MONOXIDE

U.P. Oppenheim
(Technion, Israel Institute of Technology, Haifa)


and

Hanna Goldring
(The Weizmann Institute of Science, Rehovoth, Israel)

TECHNICAL (SCIENTIFIC) NOTE No. 4

CONTRACT No. AF 61(052)-388

December, 1962



This work was supported in part by the Cambridge Research Laboratories
of the Office of Aerospace Research, United States Air Force, through
its European Office.

404042

CATALOGED BY ASTIA
AS AD NO.

404042

The Strengths of Absorption Lines in the 2-0 Band
of Carbon Monoxide

U.P. Oppenheim
(Technion, Israel Institute of Technology, Haifa)

and

Hanna Goldring
(The Weizmann Institute of Science, Rehovoth, Israel)

A new determination of the strengths of absorption lines in the 2-0 band of CO has been carried out. Very recently the strengths of lines in this band were reported by Kostkowski and Bass⁽¹⁾, and also briefly by Benedict et al.⁽²⁾. The present observations were made with better spectral resolution than the ones made by these workers. The spectrometer was a 5-m grating instrument used in double pass.⁽³⁾ The spectral slit width used was about 0.07 cm^{-1} . The studies were made with gas at approximately atmospheric pressure, and with sample lengths varying from 1 to 30 cm. The sample temperature was 300°K .

The observed line widths were between 0.05 cm^{-1} to 0.10 cm^{-1} . Under these conditions the instrumental slit corrections⁽⁴⁾ to the strengths amount to no more than 3 per cent. Since this was within the experimental error, no instrumental corrections were applied and the strengths were obtained by direct integration. It was estimated that wing corrections amounted to one or two per cent at most and they were therefore neglected.

Using the notation of Benedict et al.⁽⁵⁾, the strength S_m^0 per unit gas pressure

of an individual line is given by

$$S_m^o = S_{2-0}^o \frac{F_m |m| \nu_m e^{-E(J)/kT}}{Q(T) \bar{\nu}} \quad (1)$$

Here $Q(T)$ is the rotational partition sum ($Q(300) = 108.86$) and $\bar{\nu}$ the effective vibrational frequency⁽⁵⁾ ($\bar{\nu}_{2-0} = 4282.1 \text{ cm}^{-1}$). The exponential factors were calculated from the accurately known^(6,7) spectroscopic constants of CO. F_m expresses the effect of the vibration-rotation interaction. It may be written for the present case, according to Herman and Wallis⁽⁸⁾, to the first order in m , as follows:

$$F_m = 1 + (0.00528 - 0.0075 \theta)m$$

where $\theta = M_0/M_1 r_e$; M_0 and M_1 are the first two coefficients of a Taylor-series expansion about the equilibrium internuclear distance r_e of the dipole moment function

$$M = M_0 + M_1(r - r_e) + \dots,$$

M_0 is almost equal to the permanent dipole moment (0.117 debye ⁽⁹⁾), and a good approximation for M_1 may be calculated from the strength of the 1-0 band⁽²⁾ ($214.8 \text{ cm}^{-2} \text{ atm}^{-1}$). The value of r_e is obtained from the rotational constant⁽⁷⁾, B . The resulting approximation to the absolute value of θ is then 0.036. Without loss of accuracy, θ has been taken equal to zero, and the vibration-rotation interaction

term may then be written as

$$F_m \sim 1 + 0.00528m.$$

Fig. 1 shows the line strengths of the present work together with those obtained by Kostkowski and Bass⁽¹⁾. The value of the band strength S_{2-o}^o which gives the best fit to the experimental points was found to be $1.51 \text{ cm}^{-2} \text{ atm}^{-1}$ at 300°K . The curve corresponding to equation (1) with $S_{2-o}^o = 1.51 \text{ cm}^{-2} \text{ atm}^{-1}$ is also shown. It is seen that the line strengths obtained in the present work are lower than those given by Kostkowski and Bass⁽¹⁾, whose measurements correspond to a band strength of $1.77 \text{ cm}^{-2} \text{ atm}^{-1}$. Benedict et al.⁽²⁾ give $S_{2-o}^o = 1.5 \text{ cm}^{-2} \text{ atm}^{-1}$, in good agreement with the present value. On the other hand, Penner⁽¹⁰⁾, by a direct determination, obtained a band strength of $1.64 \text{ cm}^{-2} \text{ atm}^{-1}$.

ACKNOWLEDGEMENTS

The authors are grateful to Professor J.H. Jaffe for his help and advice throughout this work, and to Drs. Kostkowski and Bass for showing us their results prior to publication.

REFERENCES

1. H.J. Kostkowski and A.M. Bass, J.Q.S.R.T. 1, 177 (1961).
2. W.S. Benedict, R. Herman, G.E. Moore and S. Silverman,
Astrophys. J. 135, 277 (1962).
3. J.H. Jaffe, Refraction of Gases in the Infrared, Advances in
Spectroscopy, Vol. II, p. 279, Interscience Publishers, New York (1961).
4. H.J. Kostkowski and A.M. Bass, J. Opt. Soc. Amer. 46, 1060 (1956).
5. W.S. Benedict, R.H. Herman, G.E. Moore and S. Silverman,
Canad. J. Phys. 34, 850 (1956).
6. E.K. Plyler, H.C. Allen, Jr. and E.D. Tidwell, J. Res. Nat. Bur. Stand.
61, 53 (1958).
7. D.H. Rank, D.P. Eastman, B.S. Rao and T.A. Wiggins, J. Opt. Soc. Amer.
51, 929 (1961).
8. R. Herman and R. Wallis, J. Chem. Phys. 23, 637 (1955).
9. Table of Dielectric Constants and Electric Dipole Moments of Substances
in the Gaseous State, U.S. Dept. of Commerce, Nat. Bur. Stand.,
Washington, D.C. (1953).
10. S.S. Penner and D. Weber, J. Chem. Phys. 19, 807 (1951).




Fig. 1 Carbon monoxide line strengths in 2-0 band at 300°K.